



Cytokinetics

Sarcomere Directed Therapies

EMPOWERING
MUSCLE
EMPOWERING
LIVES



John, diagnosed with heart failure



Jillian, diagnosed with HCM



Chuck, diagnosed with ALS

Forward-Looking Statements

This Presentation contains forward-looking statements for purposes of the Private Securities Litigation Reform Act of 1995 (the “Act”). Cytokinetics disclaims any intent or obligation to update these forward-looking statements and claims the protection of the Act's Safe Harbor for forward-looking statements. Examples of such statements include, but are not limited to, statements related Cytokinetics' research and development and commercial readiness activities, including the initiation, conduct, design, enrollment, progress, continuation, completion, timing and results of clinical trials, projections regarding growing prevalence, low survival rates and market opportunity in heart failure, hypertrophic cardiomyopathy (HCM) or amyotrophic lateral sclerosis (ALS); projections regarding the size of the addressable patient population for *omecamtiv mecarbil*, CK-274 or *reldesemtiv*; Cytokinetics' commercial readiness for *omecamtiv mecarbil*; the likelihood of approval and timing for regulatory approval of *omecamtiv mecarbil* or any of our other drug candidates; the submission of a new drug application (NDA) to the FDA for *omecamtiv mecarbil* in 2021; the timing of commencement of COURAGE-ALS, a phase 3 clinical trial of *reldesemtiv* or the timing of commencement of a phase 3 clinical trial of CK-274; the timing of any potential commercial launch of our product candidates, if approved; commercial opportunities for our product candidates; Cytokinetics' cash runway; interactions with the FDA; the properties, potential benefits and commercial potential of CK-274, *omecamtiv mecarbil*, *reldesemtiv* and Cytokinetics' other drug candidates. Such statements are based on management's current expectations; but actual results may differ materially due to various risks and uncertainties, including, but not limited to, potential difficulties or delays in the development, testing, regulatory approvals for trial commencement, progression or product sale or manufacturing, or production of Cytokinetics' drug candidates that could slow or prevent clinical development or product approval, including risks that current and past results of clinical trials or preclinical studies may not be indicative of future clinical trial results, patient enrollment for or conduct of clinical trials may be difficult or delayed, Cytokinetics' drug candidates may have adverse side effects or inadequate therapeutic efficacy, the FDA or foreign regulatory agencies may delay or limit Cytokinetics' ability to conduct clinical trials, and Cytokinetics may be unable to obtain or maintain patent or trade secret protection for its intellectual property; Cytokinetics may incur unanticipated research, development and other costs or be unable to obtain financing necessary to conduct development of its products; standards of care may change, rendering Cytokinetics' drug candidates obsolete; and competitive products or alternative therapies may be developed by others for the treatment of indications Cytokinetics' drug candidates and potential drug candidates may target. These forward-looking statements speak only as of the date they are made, and Cytokinetics undertakes no obligation to subsequently update any such statement, except as required by law. For further information regarding these and other risks related to Cytokinetics' business, investors should consult Cytokinetics' filings with the Securities and Exchange Commission (the “SEC”).

Sarcomere Directed Therapies

OUR MISSION

To bring forward new medicines to improve the healthspan of people with devastating cardiovascular and neuromuscular diseases of impaired muscle function.

VISION 2025

Leading with Science,
Delivering for Patients

As always, we will support disease advocacy groups elevating the patient voice and live by our values of integrity, fairness and compassion in all that we do.

Our vision is to be the leading muscle biology biopharma company that meaningfully improves the lives of patients with diseases of impaired muscle function through access to our pioneering medicines

Achieve regulatory approvals for at least two drugs arising from our pipeline

Build commercial capabilities to market and sell our medicines reflective of their innovation and value


Generate sustainable and growing revenues from product sales

Double our development pipeline to include ten therapeutic programs


Expand our discovery platform to muscle energetics, growth and metabolism

Be the science-driven company people want to join and partner with

Executing On Our Vision



Pipeline of Novel Muscle-Directed Drug Candidates



* Astellas to provide co-funding in exchange for low single-digit royalty
 All drug candidates above are investigational products and are not approved as safe or effective for any indication.

Sarcomere Directed Drug Development

CARDIAC MUSCLE

Omecamtiv Mecarbil

CK-136 (AMG 594)


CK-274, CK-271

Omeamtiv ***Mecarbil***

Contractile Dysfunction Underlies Heart Failure

Increased / Preserved Cardiac Contractility

- Non-obstructive Hypertrophic Cardiomyopathy (nHCM)
- **Obstructive Hypertrophic Cardiomyopathy (oHCM)**
- Heart Failure with Preserved Ejection Fraction (certain HFpEF subsets)




Decreased Cardiac Contractility

- **Heart Failure with Reduced Ejection Fraction (HFrEF)**
- Genetic Dilated Cardiomyopathy
- Pulmonary Hypertension with Right Ventricular Heart Failure


Heart Failure: Growing Prevalence and High Readmission Rates

6 million people have heart failure in the United States

Prevalence Expected to Increase by 46% from 2012 – 2030



Mozzafarian, et al. *Circulation* 2016; 133: e38-360



1. Adams et al. *Am Heart J* 2006; 149:209-16
 2. Chen et al. *JAMA* 2011;306:1669-78
 3. Dickstein et al. *Eur Heart J* 2008;29:2388-442
 4. Korda, et al. *BMC Health Serv Res.* 2017;21;17(1):220.
 5. Krumholz et al. *Arch Intern Med* 1997;157:99 – 105


6. Krumholz et al. *Circ Cardiovasc Qual Outcomes* 2009;2(5):407-13
 7. Loehr et al. *Am J Cardiol* 2008;101:1016-22
 8. Whellan et al. *Circulation* 2010 Jan;3(1):33-40

High Economic Burden of Heart Failure

Heart failure costs ~\$123 billion annually, representing 33% of total Medicare budget^{1,2}

Heart failure is the most frequent diagnosis for hospitalized Medicare patients in the US^{1,2}

Inpatient Admission Rates for HF Patients 6X Higher than Non-HF Patients¹



1. Milliman Analysis of Medicare 5% Sample 2011-2012 (2012 index year, 2011 look back year)

2. Milliman Analysis of Medicare 5% Sample (2014 index year, 2013 look back year) and Office of the Actuary 2016 Board of Trustees Report. The costs only include Part A & B costs

Significant Unmet Need in HFrEF

Proprietary market research suggests need for novel therapy



Market research suggests need for novel therapy

Physicians say newly approved therapies have prolonged survival, decreased hospital visits, but still **see need for other therapies that reduce mortality**




Drugs that do not affect renal function

Most physicians recognize negative effect therapies such as aldosterone antagonists have **on renal function**



Drugs that do not affect BP

BP often limiting factor for up titration and therapy initiation
Need efficacious drugs **that do not result in hypotension**



Drugs that enhance cardiac performance

Need drugs that target **novel/more specific molecular targets**
Need targets other than the neurohormonal pathway



Disease modifying therapies

Need drugs that safely enhance contractility
Increased EF most frequently mentioned desired measure




Drugs that increase QoL


Patient management will improve **with drugs that increase QoL**
Patient QoL decreases as they lose the ability to perform daily tasks

Sarcomere Directed Drug Development

Cardiac muscle



The sarcomere is a molecular structure found in skeletal and cardiac muscle that enables cardiac myocytes to contract and generate force



Pivotal Phase 3 Trial Design



Landmark clinical trial results published in NEJM

Overview

Enrolled 8,256 patients at ~1,000 sites in 35 countries

Primary Endpoint

Composite of time to cardiovascular (CV) death or first HF event*, whichever occurs first

Secondary Endpoints

- Time to CV death
- Change in Kansas City Cardiomyopathy Questionnaire Total Symptoms Score (KCCQ TSS) from baseline to Week 24
- Time to first HF hospitalization
- Time to all-cause death

Key Design Points

- Dose optimization based on trough concentration of *omecamtiv mecarbil* at 2 weeks and 6 weeks
- High risk patients enrolled from inpatient and outpatient settings
- Designed to provide 90% statistical power to assess risk of CV death

*An HF event defined as the presentation of the subject for an urgent, unscheduled clinic/office/ED visit, or hospital admission, with a primary diagnosis of HF, where the patient exhibits new or worsening symptoms of HF on presentation, has objective evidence of new or worsening HF, and receives initiation or intensification of treatment specifically for HF (Hicks et al, 2015). Changes to oral diuretic therapy do not qualify as initiation or intensification of treatment.

Baseline Characteristics




Characteristic	OM (N=4120)	Placebo (N=4112)
<i>Demographics</i>		
Age (years), median (Q1, Q3)	66 (58, 73)	66 (58, 73)
Sex, female, n (%)	875 (21.2)	874 (21.3)
White/Asian/Black/other, %	78/9/7/7	78/9/7/7
<i>Heart Failure History and Medical Conditions</i>		
LVEF (%), mean (SD)	26.6 (6.3)	26.5 (6.3)
NYHA class, II/III/IV, %	53/44/3	53/44/3
Ischemic etiology, %	53.2	54.0
Atrial fib/flutter at screening, %	27.8	26.7
Type 2 diabetes, %	40.1	40.3

Characteristic	OM (N=4120)	Placebo (N=4112)
<i>Vitals and Laboratory Parameters</i>		
NT-proBNP (pg/mL), median (Q1, Q3)	1977 (980, 4061)	2025 (1000, 4105)
SBP (mmHg), mean (SD)	116 (15)	117 (15)
Heart rate, mean (SD)	72 (12)	72 (12)
eGFR (mL/min/1.73m ²), median (Q1, Q3)	59 (44, 74)	59 (44, 74)
Cardiac TnI (ng/mL), median (Q3)	0.027 (0.052)	0.027 (0.052)
<i>Medications and Cardiac Devices</i>		
ACEI/ARB/ARNi, %	87	87
ARNi, %	20	19
BB, %	94	94
MRA, %	78	78
SGLT2i, %	2.5	2.8
CRT, %	14	14
ICD, %	32	31

ACEi, angiotensin-converting enzyme inhibitor; ARB, angiotensin receptor blocker; ARNi, angiotensin receptor-neprilysin inhibitor; BB, beta blocker; CRT, cardiac resynchronization therapy; eGFR, estimated glomerular filtration rate; fib, fibrillation; hsTnI, high-sensitivity troponin I; ICD, implantable cardioverter-defibrillator; LVEF, left ventricular ejection fraction; MRA, mineralocorticoid receptor antagonist; NT-proBNP, N-terminal pro-B-type natriuretic peptide; NYHA, New York Heart Association; Q, quartile; SBP, systolic blood pressure; SGLT2i, sodium-glucose co-transporter 2 inhibitor.

Primary Composite Endpoint


Time to first HF event or CV death



HR = 0.92
(95% CI, 0.86-0.99)

P = 0.025

	0	6	12	18	24	30	36
Patients at risk, n							
Placebo	4112	3310	2889	2102	1349	647	141
OM	4120	3391	2953	2158	1430	700	164




Primary Composite Components and KCCQ TSS



CV Death

HR = 1.01 (95% CI, 0.92-1.11)


P = 0.86




Heart Failure Event

HR = 0.93 (95% CI, 0.86-1.00)

P = 0.063



Change in KCCQ TSS from Baseline to Week 24



No reduction in the secondary endpoint of time to CV death was observed

Laboratory and Safety Events




Variable	<i>Omeamtiv Mecarbil</i> (N=4110)	Placebo (N=4101)	Relative Risk or Difference (95% CI)
<i>Laboratory value change from baseline to Week 24</i>			
Systolic blood pressure - mmHg, mean (SD)	1.4 (15.3)	1.5 (15.6)	-0.1 (-0.9, 0.6)
Heart rate, bpm, mean (SD)	-2.1 (12.6)	-0.5 (12.8)	-1.6 (-2.2, -1.0)
Cardiac Troponin I, ng/L, median (Q1, Q3)	0.004 (-0.002, 0.021)	0.000 (-0.009, 0.008)	0.004 (0.003, 0.005)
NT-proBNP, pg/mL, median (Q1, Q3)	-251 (-1180, 295)	-180 (-915, 441)	0.90 (0.86, 0.94)
<i>Adverse events (AEs)</i>			
Any serious AE, n (%)	2373 (57.7)	2435 (59.4)	0.97 (0.94, 1.01)
Drug discontinuation due to AE, n (%)	371 (9.0)	382 (9.3)	0.97 (0.85, 1.11)
Adverse events of interest			
Ventricular tachyarrhythmias	290 (7.1)	304 (7.4)	0.95 (0.82, 1.11)
Torsade de pointes/QT prolongation	176 (4.3)	195 (4.8)	0.90 (0.74, 1.10)
SAE of ventricular arrhythmia requiring treatment	119 (2.9)	127 (3.1)	0.93 (0.73, 1.20)
Adjudicated major cardiac ischemic events, n (%)	200 (4.9)	188 (4.6)	1.06 (0.87, 1.29)
Myocardial infarction	122 (3.0)	118 (2.9)	
Hospitalized for unstable angina	25 (0.6)	12 (0.3)	
Coronary revascularization	115 (2.8)	117 (2.9)	
Adjudicated Strokes	76 (1.8)	112 (2.7)	0.68 (0.51, 0.91)

Treatment Effect Increased Progressively As Baseline EF Decreased


In EF ≤22%, 11.8 needed-to-treat to prevent 1 event over 3 years



Incidence of Primary Composite Endpoint



Relative Treatment Effect on Primary Composite Endpoint



Greater Treatment Effect in More Severe HF



Results of the primary outcome in pre-specified subgroups showed greater treatment effect in patients with markers of more severe heart failure, including patients with LVEF $\leq 28\%$: (n=4,456) HR 0.84; 95% CI 0.77, 0.92


Subgroup	No. of Events/ No. of Patients	Hazard Ratio (95% CI)	Norm p-value	ARR
All Patients	3103/8232	0.92 (0.86, 0.99)	0.025	2.1%
LVEF $\leq 28\%$	1821/4456	0.84 (0.77, 0.92)	<0.001	4.9%
Outpatients	1255/3304	0.83 (0.75, 0.93)	0.001	5.0%
Inpatients	566/1152	0.86 (0.73, 1.02)	0.084	3.9%
Hosp <3 mos	1200/2688	0.83 (0.74, 0.93)	0.001	5.2%
Class III/IV	1055/2132	0.80 (0.71, 0.90)	<0.001	7.0%
NT-proBNP >2000	1249/2431	0.77 (0.69, 0.87)	<0.001	8.1%
SBP <110	843/1820	0.81 (0.70, 0.92)	0.002	7.4%

Increased Treatment Effect with Severe HF

Severe HF defined as NYHA III-IV, EF \leq 30%, HF hospitalization in last 6 months



Heart Failure
World Congress on
Acute Heart Failure
2021



Treatment effect for primary endpoint in severe HF

HR = 0.80 (0.71, 0.90)

Absolute risk reduction 8.3 events/100 pt-years

NNT = 12


Source: Felker GM, Omecamtiv Mecarbil in Patients with Severe Heart Failure: An Analysis from GALACTIC-HF, ESC Heart Failure 2021, June 2021

Increased Treatment Effect with Higher NT-proBNP




Heart Failure
 & World Congress on
 Acute Heart Failure 2021

Primary Composite Outcome




HF Hospitalization



Source: McMurray JM, Efficacy of omecamtiv mecarbil in HFrEF according to NT-proBNP level: Insights from the GALACTIC-HF trial, ESC Heart Failure 2021, June 2021

Large Number of Patients At Potential US Launch Of *Omecamtiv Mecarbil*



1.2 – 2.0M patients at potential launch

1) National Center for Health Statistics. National Health and Nutrition Examination Survey (NHANES) as accessed 4/1/2019 at website. <https://www.cdc.gov/nchs/nhanes/>. – data from 2013-2016 as quotes in Benjamin 2019 Circulation. 2019;139:e56–e528.

DOI: 10.1161/

2) EF based on distribution as presented in Dunlay et al Circ Heart Fail. 2012;5:720-726,

3) 2.1% annual growth rate:1.9% annual growth rate of patient population 65+ (UN World Populations Prospects Nov 2019) and a 0.2% mortality impact of HF treatment (doi: 10.1136/bmj.l223 | BMJ 2019;364:l223)


Potential to Offset Medicare Hospitalization Costs

Outcomes from GALACTIC-HF may translate into economic benefits to payers and IDNs

Hospitalization drives cost for Medicare patients¹

- Mean cost per HFrEF hospitalization: **\$10,735**
- Mean cost for 30-day post-hospitalization care: **\$7,060**
- **Total 30-day cost for HFrEF hospitalization & post-hospitalization care: \$17,795**

Patients on *omecamtiv mecarbil* showed reductions in both hospital admissions and re-admissions²




1. Desai et al, Yale University School of Medicine, AHA 2020; Congest Heart Fail. 2011 Jul-Aug; 17(4): 10.1111/j.1751-7133.2011.00246.x.

2. GALACTIC-HF

Fit-for-Purpose Sales Team: Face-to-Face & Virtual Visits

Illustration




% of Face-to-Face Visits		
Heavy face-to-face	Mix of face-to-face and remote	Minimum face-to-face
Engagement Description		
Similar to traditional engagement – rep spends most of the time in face-to-face interaction	Hybrid engagement – mix of face-to-face and virtual visits to sequence interactions depending on customer needs and constraints. Remote resources deployed (i.e., samples, speakers, literature)	Dominant use of virtual platforms. Interaction is primarily over scheduled virtual visits or phone calls in response to office queries. Remote resources deployed (i.e., samples, speakers, literature)

Note: Sep'20 Access Monitor stats indicate the growing preference for face-face visits. Based on Access Monitor and Voice of Patient & Provider surveys


Applied Analytics Will Inform Channel Mix and Deployment

Patient and HCP Heat Map in HFReF



Deploy to Hot Spots

Physician Engagement Type by Geography



Note: Based on 2020 cycle 1 AffinityMonitor™ metrics for LHMs; LHM engagement was considered to be the average engagement of rated HCPs within each LHMs; LHMs are ZS designed market which are homogeneous market within LHM boundaries

Second Phase 3 Clinical Trial Underway

Investigating effect of *omecamtiv mecarbil* on exercise tolerance

Enrollment complete; results expected in 2H 2022

Primary Endpoint
Change in peak VO ₂ on CPET from baseline to Week 20
Second Endpoints
<ul style="list-style-type: none"> Change in total workload during CPET from baseline to Week 20 Change in ventilatory efficiency (VE/VCO₂ slope) during CPET from baseline to Week 20 Change in average daily activity units measured over 2 weeks from baseline to Week 18-20 by accelerometry


Study Plan	
Total Countries Planned	9
Active Countries	4
Total Sites Planned	92
Activated Sites	69
Total Patients Planned	270

Key Design Points

- Designed to enroll approximately 270 patients
- 90% power
- Patients must have LVEF ≤35 percent, be NYHA heart failure class II or III, and have reduced exercise capacity
- Patients randomized 2:1 to *omecamtiv mecarbil*

VO₂ = Oxygen Uptake; CPET = Cardio-Pulmonary Exercise Testing; VE = Ventilatory Efficiency


Clinical Trial Overview



*Screening echocardiogram is not required if an appropriate LVEF assessment has been performed within one year

CK-274

Symptomatic HCM: Orphan Indication



Source: #26 SHA 2016-2021 Patient Claims Data; #20 Cogent HC 2020 DoF

Significant Unmet Need in HCM

Current therapies do not target underlying disease



HCM is an inherited cardiovascular disease

1 in 500 have genetic mutation
1 in 3200 have HCM
Subset of patients have progressive symptoms, atrial fibrillation, stroke, sudden death



Surgical intervention not permanent solution

Invasive therapy to reduce septal thickness is effective
Surgical myectomy or percutaneous ablation



Current medical therapy does not target underlying disease

Indirect mechanisms of action with systemic side effects
Variable efficacy, often inadequate

CK-274: Next-In-Class Cardiac Myosin Inhibitor

Potential treatment for patients with HCM




- Selective allosteric inhibitor of cardiac myosin discovered by company scientists independent of collaborations
- Potential *in vivo* pharmacodynamic advantages related to distinctive binding site
- Optimized for
 - Onset of action (reach steady state within two weeks)
 - Rapid reversibility of effect
 - Minimal drug-drug interactions
 - Favorable tolerability
 - Ease of titration for personalized dosing
- Clear pharmacokinetic/pharmacodynamic (PK/PD) relationship observed
- Shallow exposure-response relationship

SAD & MAD Results Support Progression to Phase 2


Preclinical data translated to healthy participants

MAD PK: Steady-State Achieved After 14 Days of Dosing



Shallow Exposure-Response Relationship Observed Pre-clinically Appears to Have Translated to Humans, May Enable Flexible Dose Optimization in Humans

PK/PD Relationship of CK-274 for Ejection Fraction (LVEF)




Graphs show LVEF as a function of exposure; data points represent observed values in dogs and humans.

Decrease in LVEF as function of exposure is similar in humans and dogs.

Phase 2 Clinical Trial Design



Two sequential dose-finding cohorts (with third cohort assessing patients on *disopyramide*)



	Dose 1	Dose 2	Dose 3
Cohort 1	5 mg	10 mg	15 mg
Cohort 2	10 mg	20 mg	30 mg

Patient Enrollment and Dosing



41 Total Enrolled Patients


Final Dose Achieved (N)						
	Cohort 1			Cohort 2		
Placebo	5 mg	10 mg	15 mg	10 mg	20 mg	30 mg
13	4	5	5	9	4	1

Baseline Echocardiographic Data




Characteristic, mean	Baseline (Day 1 Pre-dose)		
	Placebo C1 + C2 Combined (N = 13)	CK-274	
		Cohort 1 (N = 14)	Cohort 2 (N = 14)
LVEF (%)	74.5	73.2	75.4
LVOT-G, Rest (mmHg)	52.1	53.8	58.2
LVOT-G, Valsalva (mmHg)	84.6	74.4	82.3

High Response Rates on Treatment with CK-274




Responder Definition:
Resting LVOT-G <30 mmHg
and post-Valsalva LVOT-G
<50 mmHg at Week 10

Resting Left Ventricular Outflow Tract Gradient



Mean \pm SEM	Valsalva LVOT-G (mmHg)				
	Baseline	Week 2	Week 4	Week 6	Week 10
Placebo (n=13)	52.1	45.0	47.1	49.0	44.0
Cohort 1 (n = 14)	53.8	24.3	27.3	13.9	13.4
p-value vs placebo	-	0.007	0.025	<0.0001	0.0003
Cohort 2 (n = 14)	58.2	15.5	16.1	10.9	15.1
p-value vs placebo	-	0.0002	0.0006	<0.0001	0.0004

Post-Valsalva Left Ventricular Outflow Tract Gradient



Mean ± SEM	Valsalva LVOT-G (mmHg)				
	Baseline	Week 2	Week 4	Week 6	Week 10
Placebo (n=13)	84.6	71.3	71.3	73.4	76
Cohort 1 (n = 14)	74.4	51.3	46.1	37.1	38.1
p-value vs placebo	-	0.097	0.038	0.0003	0.001
Cohort 2 (n = 14)	82.3	32.3	31.5	30.3	29.8
p-value vs placebo	-	0.0005	0.0005	<0.0001	<0.0001

Safety Data



- Incidence of adverse events on CK-274 similar to placebo and mild or moderate
- There were no treatment related serious adverse events reported by investigators
- No patients who received CK-274 in Cohort 1 had an LVEF <50%
- In Cohort 2, one patient with LVEF at baseline of 58% was up titrated to 20 mg and experienced transient LVEF reduction to <50% (remaining above 40%) requiring down titration
- No interruptions or discontinuations of treatment with CK-274 occurred across both cohorts

Open Label Extension Trial



REDWOOD-HCM OLE open for eligible patients who completed REDWOOD-HCM


- Primary endpoint: incidence of AEs & LVEF <50
- Secondary endpoints: measures of long-term effects of CK-274 on LVOT-G; assessments of steady-state pharmacokinetics.
 - Cardiac MRI sub-study to assess changes in cardiac morphology, function and fibrosis
- Individually optimized dose starts at lowest dose in prespecified range with echo-guided dose titration
- Initial dose and highest target dose informed by interim analyses from REDWOOD-HCM

OLE: Escalating doses based on echo-guided dose titration

CK-274: Clinical Development Plan for HCM


Engaging regulatory authorities to inform Phase 3

Type C meeting with FDA to review Phase 3 design; End-of-Phase 2 meeting to review dose selection rationale for Phase 3



Novel Approach May Address Multiple Unmet Patient Needs

No FDA-approved therapies



CK-274: Collaborations & Agreements

RTW Investments, LP & Ji Xing Pharmaceuticals Limited



RTW & Ji Xing Pharma Licensing Collaboration, Funding Commitments & Royalty Monetization

RTW Investments committed capital, funding and sale proceeds of \$250M to Cytokinetics

Ji Xing Pharma to develop & commercialize CK-274 in China, subject to royalties and up to \$200M in milestone payments

RTW Investments purchased equity and royalty; provides access to capital for development of CK-274

Ji Xing Pharma

Ji Xing to develop & commercialize CK-274 in Greater China and Taiwan

Cytokinetics receives **\$25M upfront**; eligible to receive **\$200M** in development & commercial milestones & double-digit royalties on sales of CK-274 in licensed territory

RTW: Funding for Development of CK-274

Cytokinetics receives options for additional funding for further development of CK-274 in HCMs:

- Eligible for **\$45M** in each of 2 tranches (upon initiation of global registration programs in oHCM and nHCM) in exchange for 2% royalty on sales in U.S. & certain European countries
- If **full \$90M** received, Cytokinetics pays RTW 4% royalty on sales of CK-274 in U.S. & certain European countries, subject to royalty reductions for potential other indications

RTW: Other Purchases

RTW purchased Cytokinetics' royalty rights **on future sales of mavacamten** for **\$85M**

RTW purchased **\$50M of Cytokinetics' common stock** at \$25 per share

Building Synergistic Commercial Capabilities

Building Today...

Building commercial organization focused on hospitalized CV patients and HCPs to optimize opportunity for *omecamtiv mecarbil*

- Cultivate advocacy with CV patients and HCPs

To Lead Tomorrow

Establish Cytokinetics as a CV leader by leveraging commercial capabilities for future product launches

- Significant overlap between HFrEF & HCM accounts
- Simultaneously gain experience in HFrEF & HCM



6,000+

Hospitals and
CoEs in US



1,100

Highest Value
Hospitals & CoEs



~75% HFrEF Patients

~78% HCM Patients

IQVIA HPD - Q3'18 - Q2'19

Sarcomere Directed Drug Development

SKELETAL MUSCLE


Reldesemtiv

CK-601

Sarcomere Directed Drug Development

Skeletal muscle

The sarcomere is a molecular structure found in skeletal and cardiac muscle that enables skeletal myocytes to contract and generate force



Reldesemtiv

Phase 2 Clinical Trial in ALS


Results presented at American Academy of Neurology 2019

Parallel group, dose ranging study enrolled 458 patients with ALS in the US, Canada, Australia and Europe evaluating change from baseline in the percent predicted slow vital capacity (SVC) at 12 weeks of treatment with *reldesemtiv* or placebo



Primary Endpoint: SVC

Change from baseline in percent predicted SVC at week 12




Primary Analysis*
P = 0.11
for weighted
dose-response
relationship

*Based on Mixed Model for Repeated Measures (MMRM) with the contrasts of (-5, -1, 3, 3) for placebo, *reldesemtiv* 150 mg, 300 mg and 450 mg BID, respectively


Change From Baseline: All Active vs Placebo*

Results support progression to potential Phase 3 clinical trial

SVC Change From Baseline (All Active vs Placebo)




ALSFRS-R Change From Baseline (All Active vs Placebo)




*post hoc analysis
FORTITUDE-ALS did not achieve statistical significance, but patients on all dose groups of *reldecentiv* declined less than patients on placebo

Subgroup Analyses*

Percent Predicted SVC




ALSFERS-R Total Score



*FORTITUDE-ALS did not achieve statistical significance, but patients on all dose groups of *reldesemtiv* declined less than patients on placebo


Post-Hoc Analyses Inform Potential Path Forward

Change From Baseline in ALSFRS-R by Progressor Tertiles



Probability of No New DME* Over Time With Treatment With *Reldesemtiv*


DME (Durable Medical Equipment): Manual wheelchair, power wheelchair, NIV, Augmentative Language Device, PEG



Planned Phase 3 Clinical Trial Design

Trial to open for enrollment in 2021

Enrolling 555 patients with ALS in the US, Canada, Australia and Europe evaluating change from baseline ALSFRS-R at 24 weeks of treatment with *reldesemtiv* or placebo



Study Visits	Screen	D1	W2	W4	W8	W12	W16	W20	W24	W26	W28	W32	W36	W40	W44	W48	W52 FU
ALSFRS-R	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑
FVC	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑
Lab	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑
Muscle Strength	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑

↑ In-Clinic ↑ Remote ↑ Both In-Clinic & Remote

Sarcomere Directed Therapies

CORPORATE PROFILE

Robust Pipeline, Solid Financial Position

Pipeline*	1 Positive trial readout in 2021	2 Pivotal trials in 2021	3 Potential FDA approvals by 2025	5 Clinical stage programs	10 Development programs by 2025
Programs*	<p>Heart Failure <i>Omecamtiv mecarbil</i></p> <ul style="list-style-type: none"> Positive trial results from GALACTIC-HF Phase 3 exercise capacity trial results early 2022 <p>CK-136</p> <ul style="list-style-type: none"> Phase 1 	<p>HCM <i>CK-274</i></p> <ul style="list-style-type: none"> Positive results from REDWOOD-HCM Expect to begin Phase 3 trial by year end 	<p>ALS <i>Reldesemtiv</i></p> <ul style="list-style-type: none"> Expect to begin COURAGE-ALS, Phase 3 trial, in Q3 2021 	<p>Ongoing R&D</p> <p>Additional research in muscle biology, energetics & metabolism</p> 	
Foundations	 <p>207 Full time employees</p>		<p>\$460M* At Q1 2021</p>		

* In July 2021, Cytokinetics raised \$275 million through a public offering of common stock. Timelines and milestones reflect Cytokinetics' current expectations and beliefs

Cytokinetics Financing History

As of 3/31/2021, with estimated proceeds from 7/23/21 offering

in millions

	As of 3/31/2021	Financing	Equity	Upfront Cash, Option, & Milestones Reimbursement	R&D	Total
Investors	Private Investors (VCs)		\$116			\$116
	IPO		\$94			\$94
	Public Post-IPO/Other		\$906			\$906
	Term Loan	\$45				\$45
	Convertible Debt (net)*	\$120.5				\$120.5
			\$165.5	\$1,116		
Strategic Partners & Grants	RTW/Ji Xing		\$50	\$113		\$163
	Astellas		\$10	\$130	\$103	\$243
	Amgen		\$43	\$145	\$58	\$246
	Royalty Pharma		\$10	\$90	-	\$100
	GSK		\$24	\$22	\$33	\$79
	AstraZeneca		-	-	\$2	\$2
	MyoKardia		-	-	\$2	\$2
	Global Blood		-	-	\$2	\$2
	Grants (ALS Assoc/NINDS/other)		-	\$6	-	\$6
				\$137	\$506	\$200

Capital raised:
combination of
strategic partners
and investors

*Net of fees and expenses, and Capped Call costs

Balance Sheet & Financial Guidance

Ended Q1 with 2+ years cash runway based on 2021 guidance

2021 Condensed Balance Sheet

As of 3/31/2021

	<i>in millions</i>
	Total
Cash and investments	\$460.2
Leased assets	\$86.1
Other assets	\$30.8
Total Assets	\$577.1
Debt	\$134.0
Liability related to sale of future royalties	\$168.9
Deferred Revenue	\$87.0
Lease liability	\$85.6
Other liabilities	\$33.7
Total Liabilities	\$509.2
Working capital	\$397.2
Accumulated deficit	(\$1,039.4)
Stockholders' equity	\$67.8
Wtd Avg Basic Shares Outstanding	71.2

2021 Financial Guidance

	<i>in millions</i>
	Total
Cash Revenue	\$23 – 28
Cash Operating Expenses*	\$230 – 250
Net	~\$195-215

*We expect to revise our financial guidance mid-year once we finalize strategies and potential commercial launch plans for *omecamtiv mecarbil*. Executing on those strategies and plans may result in our incurring significant additional expenses that were not included in our current financial guidance.

Expected Upcoming 2021 Milestones

Expect to Begin **Phase 3 Trial of CK-274** by Year End

Submit US NDA for **omecamtiv mecarbil** in 2H 2021

Expect results from **METEORIC-HF** in early 2022

Start COURAGE-ALS, Phase 3 Clinical Trial of **Reldesemtiv** in Patients with ALS, in Q3



Cytokinetics

Sarcomere Directed Therapies

**THANK
YOU**



John, diagnosed with heart failure



Jillian, diagnosed with HCM



Chuck, diagnosed with ALS